



Access fun Grade 8–12 quizzes, matric past papers, K53 learner mock tests, and NBT prep!

All in one easy-to-use app.

DOWNLOAD GO STUDY NOW



Tap on the buttons above to download the app

 www.gostudy.club



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**SENIOR CERTIFICATE EXAMINATIONS/
SENIORSERTIFIKAAT-EKSAMEN
NATIONAL SENIOR CERTIFICATE EXAMINATIONS/
NASIONALE SENIORSERTIFIKAAT-EKSAMEN**

**MATHEMATICS P2/
WISKUNDE V2**

MARKING GUIDELINES/NASIENRIGLYNE

2021

**MARKS: 150
PUNTE: 150**

**These marking guidelines consist of 23 pages.
*Hierdie nasienriglyne bestaan uit 23 bladsye.***

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.

LET WEL:

- As 'n kandidaat 'n vraag TWEE KEER beantwoord, sien slegs die EERSTE poging na.
- As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, sien die doodgetrekte poging na.
- Volgehoue akkuraatheid word in ALLE aspekte van die memorandum toegepas. Hou op nasien by die tweede berekeningsfout.
- Aanvaar van antwoorde/waardes om 'n probleem op te los, word NIE toegelaat nie.

| GEOMETRY | |
|----------|--|
| S | A mark for a correct statement (A statement mark is independent of a reason) |
| | <i>'n Punt vir 'n korrekte bewering</i> (<i>'n Punt vir 'n bewering is onafhanklik van die rede</i>) |
| R | A mark for the correct reason (A reason mark may only be awarded if the statement is correct) |
| | <i>'n Punt vir 'n korrekte rede</i> (<i>'n Punt word slegs vir die rede toegeken as die bewering korrek is</i>) |
| S/R | Award a mark if statement AND reason are both correct |
| | <i>Ken 'n punt toe as die bewering EN rede beide korrek is</i> |

QUESTION/VRAAG 1

1.1

| | | | | | | | | | | | | | | |
|----|----|---|----|----|----|----|----|----|----|----|----|----|----|----|
| 26 | 13 | 3 | 18 | 12 | 34 | 24 | 58 | 16 | 10 | 15 | 69 | 20 | 17 | 40 |
|----|----|---|----|----|----|----|----|----|----|----|----|----|----|----|

| | | |
|----------|--|--|
| 1.1.1(a) | $\bar{x} = \frac{375}{15}$ $\bar{x} = 25 \text{ MB}$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">Answer only: Full marks</div> | ✓ 375 ✓ answer (2) |
| 1.1.1(b) | $\sigma = 17,65 \text{ MB}$ | ✓ answer (1) |
| 1.1.2 | $25 + 17,65 = 42,65$ $\therefore 2 \text{ days}$ | ✓ 42,65 ✓ 2 (2) |
| 1.1.3 | Overall $\bar{x} = \frac{80}{100} \times 25$ $= 20 \text{ MB}$ $\frac{375 + x}{30} = 20$ $x = 600 - 375$ $= 225$ maximum total amount of data that Sam must use for the remainder of the month: 225 MB | ✓ Overall $\bar{x} = 20$ ✓ $\frac{375 + x}{30} = 20$ ✓ answer (3) |


1.2

| Wind speed in km/h (x) | 2 | 6 | 15 | 20 | 25 | 17 | 11 | 24 | 13 | 22 |
|------------------------|----|----|----|----|----|----|----|----|----|----|
| Temperature in °C (y) | 28 | 26 | 22 | 22 | 16 | 20 | 24 | 19 | 26 | 19 |

| | | |
|-------------|--|---|
| 1.2.1 | $a = 29,35$ $b = -0,46$ $\hat{y} = 29,35 - 0,46x$ | ✓ a ✓ b ✓ equation (3) |
| 1.2.2 | $y = 25,20 \text{ °C}$ (calculator) OR $\hat{y} = 29,35 - 0,46(9)$ $y = 25,21 \text{ °C}$ | ✓✓ answer (2) ✓ substitution ✓ answer (2) |
| 1.2.3 | $b < 0$, indicating that as the wind speed increases the temperature decreases. | ✓ interpretation (1) |
| [14] | | |


QUESTION/VRAAG 2

| Number of days absent | Number of learners | Cumulative frequency |
|-----------------------|--------------------|----------------------|
| $0 \leq x < 5$ | 34 | 34 |
| $5 \leq x < 10$ | 45 | 79 |
| $10 \leq x < 15$ | 98 | 177 |
| $15 \leq x < 20$ | 43 | 220 |
| $20 \leq x < 25$ | 7 | 227 |
| $25 \leq x < 30$ | 3 | 230 |

| | | |
|-----|---|--|
| 2.1 | Modal class: $10 \leq x < 15$ | ✓ answer (1) |
| 2.2 | 177 learners | ✓ answer (1) |
| 2.3 | 230 learners | ✓ answer (1) |
| 2.4 |  <p style="text-align: center;">Ogive</p> <p>Cumulative frequency</p> <p>Number of days absent</p> | <p>✓ grounding at (0; 0)</p> <p>✓ shape</p> <p>✓ upper limits</p> <p>✓ All other points correct</p> <p>(4)</p> |
| 2.5 | <p>The median is at position 115.</p> <p>□ value of median is 12 days (accept 11 – 14)</p> <div style="border: 1px solid black; padding: 5px; display: inline-block;">Answer only: Full marks</div> | <p>✓ reading off at 115</p> <p>✓ answer</p> <p>(2)</p> |

[9]

QUESTION/VRAAG 3




| | | |
|-----|--|--|
| 3.1 | $M\left(\frac{4+8}{2}; \frac{-8+0}{2}\right)$ $M(6; -4)$ | ✓ x_M ✓ y_M (2) |
| 3.2 | $m_{NS} = \frac{0 - (-16)}{8 - 0} \text{ or } m_{NQ} = \frac{0 - (-8)}{8 - 4} \text{ or } m_{QS} = \frac{-8 - (-16)}{4 - 0}$ $= 2$ | ✓ subst N and Q or N and Q or Q and S into gradient formula ✓ answer (2) |
| 3.3 | $m_{LQ} \times 2 = -1 \quad [LQ \perp NS]$ $\therefore m_{LQ} = -\frac{1}{2}$ $-8 = -\frac{1}{2}(4) + c \quad \text{OR} \quad y + 8 = -\frac{1}{2}(x - 4)$ $c = -6 \quad \quad \quad y + 8 = -\frac{1}{2}x + 2$ $\therefore y = -\frac{1}{2}x - 6$ | ✓ m_{LQ} ✓ substitution of Q ✓ calculation of c or simplification (3) |
| 3.4 | OS is the radius of circle passing through S $(x - 0)^2 + (y - 0)^2 = (16)^2$ $x^2 + y^2 = 256$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">Answer only: Full marks</div> | ✓ identifying radius = 16 ✓ Equation of circle (2) |

| | | |
|-----|--|---|
| 3.5 | $m_{RM} = m_{LQ} = -\frac{1}{2} \quad [RM \parallel LQ]$ $-4 = -\frac{1}{2}(6) + c \quad \text{OR} \quad y + 4 = -\frac{1}{2}(x - 6)$ $c = -1 \quad y + 4 = -\frac{1}{2}x + 3$ $\therefore y = -\frac{1}{2}x - 1$ <p>T(0; -1)</p> | <p>✓ m_{RM}</p> <p>✓ substitution of M(6; -4)</p> <p>✓ coordinates of T</p> <p style="text-align: right;">(3)</p> |
| 3.6 | <p>T(0; -1), P(0; -6) and S(0; -16)</p> <p>$\therefore PS = 10$ units and $TS = 15$ units</p> $\frac{LS}{RS} = \frac{PS}{TS} = \frac{2}{3} \quad \begin{array}{l} [\text{prop theorem; RM} \parallel \text{LP}] \\ \text{OR} [\text{line} \parallel \text{one side of} \\ \Delta / \text{lyn} \parallel \text{een sy v } \Delta] \end{array}$ <div style="border: 1px solid black; padding: 2px; width: fit-content; margin: 5px auto;">Answer only: Full marks</div> <p>OR</p> <p>M(6; -4), Q(4; -8) and S(0; -16)</p> <p>$MS = \sqrt{180} = 6\sqrt{5}$ and $QS = \sqrt{80} = 4\sqrt{5}$</p> $\frac{LS}{RS} = \frac{QS}{MS} = \frac{2}{3} \quad \begin{array}{l} [\text{prop theorem; RM} \parallel \text{LQ}] \\ \text{OR} [\text{line} \parallel \text{one side of} \\ \Delta / \text{lyn} \parallel \text{een sy v } \Delta] \end{array}$ <div style="border: 1px solid black; padding: 2px; width: fit-content; margin: 5px auto;">Answer only: Full marks</div> | <p>✓ PS = 10 units</p> <p>✓ TS = 15 units</p> <p>✓ answer</p> <p style="text-align: right;">(3)</p> <p>✓ $MS = 6\sqrt{5}$ units</p> <p>✓ $QS = 4\sqrt{5}$ units</p> <p>✓ answer</p> <p style="text-align: right;">(3)</p> |
| 3.7 | <p>area of PTMQ = area of ΔTSM – area of ΔPSQ</p> $= \frac{1}{2} \cdot ST \cdot \perp h_M - \frac{1}{2} \cdot PS \cdot \perp h_Q$ $= \frac{1}{2}(15)(6) - \frac{1}{2}(10)(4)$ $= 45 - 20$ $= 25 \text{ square units}$ <p>OR</p> <p>$TM = \sqrt{45} = 3\sqrt{5} = 6,71$</p> <p>$MQ = \sqrt{20} = 2\sqrt{5} = 4,47$</p> <p>$PQ = \sqrt{20} = 2\sqrt{5} = 4,47$</p> <p>area of trapezium PTMQ = $\frac{1}{2}(3\sqrt{5} + 2\sqrt{5})(2\sqrt{5})$</p> $= \frac{1}{2}(5\sqrt{5})(2\sqrt{5})$ $= 25 \text{ square units}$ | <p>✓ area of ΔTSM – area of ΔPSQ</p> <p>✓ area $\Delta TSM = 45$</p> <p>✓ area $\Delta PSQ = 20$</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> <p>✓ $TM = 3\sqrt{5}$</p> <p>✓ $MQ = 2\sqrt{5}$</p> <p>✓ $PQ = 2\sqrt{5}$</p> <p>✓ area of trapezium = $\frac{1}{2}$ (sum of sides)(height)</p> <p>✓ substitute into formula</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> |

| | | |
|--|--|---|
| | <p>OR</p> <p>$MQ = \sqrt{20} = 2\sqrt{5}$</p> <p>$PQ = \sqrt{20} = 2\sqrt{5}$</p> <p>$TP = 5$</p> <p>area of PTMQ = area of $\triangle MTP$ + area of $\triangle PQM$</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> $\text{area of PTMQ} = \frac{1}{2} TP \times \perp h_M + \frac{1}{2} MQ \times PQ$ </div> <p>area of PTMQ = $10 + 15 = 25$</p> | <p>✓ area of $\triangle MTP$ + area of $\triangle PQM$</p> <p>area of PTMQ = $\frac{1}{2}(5) \times 6 + \frac{1}{2}(2\sqrt{5})(2\sqrt{5})$</p> <p>✓ area $\triangle MTP = 10$</p> <p>✓ area $\triangle PQM = 15$</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> |
| | | [19] |

QUESTION 4



| | | |
|-----|---|---|
| 4.1 | $PV = r = \sqrt{10}$ $PV = \sqrt{(k - (-3))^2 + (1 - 4)^2} = \sqrt{10}$ $(PV)^2 = (k - (-3))^2 + (1 - 4)^2 = 10$ $k^2 + 6k + 9 + 9 = 10$ OR $(k + 3)^2 + 9 = 10$ $k^2 + 6k + 8 = 0$ $(k + 3)^2 = 1$ $(k + 4)(k + 2) = 0$ $k + 3 = 1$ or $k + 3 = -1$ $k = -4$ or $k = -2$ $\therefore k = -2$ | ✓ $PV = r = \sqrt{10}$ ✓ substitution into distance formula ✓ standard form ✓ factors ✓ answer (5) |
| 4.2 | $x^2 + 6x + y^2 - 8y + 15 = 0$ y-intercepts: $(0)^2 + 6(0) + y^2 - 8y + 15 = 0$ $(y - 3)(y - 5) = 0$ $y_C = 3$ or $y_B = 5$ $\therefore BC = 2$ units | ✓ $x = 0$ ✓ factors ✓ both values ✓ answer (4) |

| | | |
|-------|---|--|
| 4.3.1 | $m_{TC} = \frac{3-1}{0-(-2)}$ $= 1$ $\tan \alpha = 1$ $\therefore \alpha = 45^\circ$ <p>OR</p> $y = mx + 3$ $1 = m(-2) + 3$ $m_{TC} = 1$ $\tan \alpha = 1$ $\therefore \alpha = 45^\circ$ | ✓ substitution into gradient formula ✓ $\tan \alpha = 1$ ✓ answer (3) |
| 4.3.2 | $\hat{BCV} = 135^\circ$ $\therefore \hat{VWB} = 45^\circ$ <p>[ext \angle of Δ/buite \angle v Δ] [opp \angles of cyclic quad/teenoorst. \anglee v kvh] Answer only: Full marks</p> <p>OR</p> $\hat{TCO} = 45^\circ$ $\therefore \hat{VWB} = 45^\circ$ <p>[\angles of Δ/\anglee v Δ] [ext \angles of cyclic quad/buite \angle v kvh] Answer only: Full marks</p> | ✓ $\hat{BCV} = 135^\circ$ ✓ answer (2) |
| 4.4.1 | $Q(-3; -2)$ | ✓ x_Q ✓ y_Q (2) |
| 4.4.2 | $(x+3)^2 + (y+2)^2 = 10$ | ✓ LHS ✓ RHS (2) |
| 4.4.3 | $x = -2$ or $x = -4$ | ✓ $x = -2$ ✓ $x = -4$ (2) |
| | | [20] |


QUESTION/VRAAG 5

| | | |
|-------|---|--|
| 5.1 | $\tan(-x) \cdot \cos x \cdot \sin(x - 180^\circ) - 1$ $= -\tan x \cdot \cos x \cdot \sin(-(180^\circ - x)) - 1$ $= \frac{-\sin x}{\cos x} \cdot \cos x \cdot (-\sin x) - 1$ $= \sin^2 x - 1$ $= -\cos^2 x$ | $\checkmark -\tan x$ $\checkmark -\sin x \quad \checkmark \frac{-\sin x}{\cos x}$ $\checkmark \sin^2 x - 1$ $\checkmark \text{ answer}$ (5) |
| 5.2.1 | $\cos 215^\circ$ $= -\cos 35^\circ$ $= -m$ | $\checkmark \text{ reduction}$ $\checkmark \text{ answer}$ (2) |
| 5.2.2 | $\sin 20^\circ$ $= \cos 70^\circ$ $= \cos 2(35^\circ)$ $= 2\cos^2 35^\circ - 1$ $= 2m^2 - 1$ <p>OR</p> $= \sin(55^\circ - 35^\circ)$ $= \sin 55^\circ \cos 35^\circ - \cos 55^\circ \sin 35^\circ$ $= m \cdot m - \sqrt{1-m^2} \cdot \sqrt{1-m^2}$ $= m^2 - (1-m^2)$ $= 2m^2 - 1$ | $\checkmark \text{ co-function}$ $\checkmark \text{ double angle expansion}$ $\checkmark \text{ answer in terms of } m$ (3) $\checkmark \text{ compound angle expansion}$ $\checkmark \cos 55^\circ = \sqrt{1-m^2} \text{ or } \sin 35^\circ = \sqrt{1-m^2}$ $\checkmark \text{ answer in terms of } m$ (3) |
| 5.3 | $\cos 4x \cdot \cos x + \sin 4x \cdot \sin x = -0,7$ $\cos(4x - x) = -0,7$ $\text{ref } \angle = 45,57 \dots^\circ$ $3x = 180^\circ - 45,57 \dots^\circ + k \cdot 360^\circ \text{ or } 3x = 180^\circ + 45,57 \dots^\circ + k \cdot 360^\circ$ $3x = 134,43^\circ + k \cdot 360^\circ \text{ or } 3x = 225,57^\circ + k \cdot 360^\circ$ $x = 44,81^\circ + k \cdot 120^\circ; k \in \mathbb{Z} \quad x = 75,19^\circ + k \cdot 120^\circ; k \in \mathbb{Z}$ | $\checkmark \text{ compound angle}$ $\checkmark 3x = 134,43^\circ \text{ or } 225,57^\circ$ $\checkmark x = 44,81^\circ \text{ or } 75,19^\circ$ $\checkmark + k \cdot 120^\circ; k \in \mathbb{Z}$ (4) |

| | | |
|-----|--|--|
| 5.4 | $\text{RHS} = \cos^2 x - \sin^2 x$ $\text{LHS} = \frac{\sin 4x \cdot \cos 2x - 2 \cos 4x \cdot \sin x \cdot \cos x}{\tan 2x}$ $= \frac{\sin 4x \cdot \cos 2x - \cos 4x \cdot \sin 2x}{\frac{\sin 2x}{\cos 2x}}$ $= \sin(4x - 2x) \left(\frac{\cos 2x}{\sin 2x} \right)$ $= \cos 2x$ $= \cos^2 x - \sin^2 x$ $\text{LHS} = \text{RHS}$ | $\checkmark \sin 2x$ $\checkmark \frac{\sin 2x}{\cos 2x}$ $\checkmark \sin(4x - 2x)$ $\checkmark \cos 2x$ |
| | | (4) |
| | | [18] |




6.2



| | | |
|-------------|---|---|
| 6.2.1 | $A(-150^\circ; 0,5)$ $B(-30^\circ; 0,5)$ $AB = -30^\circ - (-150^\circ)$ $AB = 120^\circ$ <div>Answer only: Full marks</div> | $\checkmark AB = -30^\circ - (-150^\circ)$ \checkmark answer (2) |
| 6.2.2 | $x \in (0^\circ; 90^\circ)$ or $x \in (90^\circ; 180^\circ)$ OR $0^\circ < x < 90^\circ$ or $90^\circ < x < 180^\circ$ | $\checkmark x \in (0^\circ; 90^\circ)$ $\checkmark x \in (90^\circ; 180^\circ)$ (2) $\checkmark 0^\circ < x < 90^\circ$ $\checkmark 90^\circ < x < 180^\circ$ (2) |
| 6.2.3 | $\cos 2x = k - 3$ $k - 3 < -1$ or $k - 3 > 1$ $k < 2$ or $k > 4$ OR $k < 2$ or $k > 4$ <div>Answer only: Full marks</div> | $\checkmark k - 3 < -1$ or $k - 3 > 1$ $\checkmark k < 2$ $\checkmark k > 4$ (3) \checkmark graph of $y = \cos 2x + 3$ $\checkmark k < 2$ $\checkmark k > 4$ (3) |
| [13] | | |

QUESTION/VRAAG 7



| | | |
|-----|--|--|
| 7.1 | <p>In $\triangle BCE$:</p> $\frac{CE}{\sin \hat{B}} = \frac{BC}{\sin \hat{E}}$ $\frac{CE}{\sin 30^\circ} = \frac{BC}{\sin 2x}$ $CE = \frac{BC \sin 30^\circ}{\sin 2x}$ <p>In $\triangle CDE$:</p> $\frac{DC}{CE} = \tan \hat{E}$ $DC = \frac{BC \cdot \sin 30^\circ}{\sin 2x} (\tan x)$ $DC = \frac{BC}{4 \sin x \cos x} \left(\frac{\sin x}{\cos x} \right)$ $DC = \frac{BC}{4 \cos^2 x}$ | <p>✓ correct use of sine rule</p> <p>✓ $CE = \frac{BC \sin 30^\circ}{\sin 2x}$</p> <p>✓ correct trig ratio</p> <p>✓ Subst CE</p> <p>✓ $2 \sin x \cos x$ ✓ $\frac{\sin x}{\cos x}$</p> <p>(6)</p> |
|-----|--|--|


| | | |
|-----|--|---|
| 7.2 | $DC = \frac{BC}{4 \cos^2 30^\circ}$ $= \frac{BC}{4 \left(\frac{\sqrt{3}}{2} \right)^2}$ $= \frac{BC}{3}$ $\therefore BC = 3DC$ <p>But $AB = DC$ [opp sides of rectangle/teenoorst. sye v reghoek]</p> $\therefore BC = 3AB$ <p>Area of rectangle = $(AB)(BC)$ $= (AB)(3AB)$ $= 3AB^2$</p> | <p>✓ $DC = \frac{BC}{3}$</p> <p>✓ $BC = 3AB$</p> <p>✓ substitution into area formula</p> <p>(3)</p> |
| [9] | | |

| | | |
|------|--|---|
| | <p>OR</p> <p>$\hat{C}_3 = 3x$ [ext \angle of cyclic quad/<i>buite</i> \angle v <i>kvh</i>]</p> <p>$\hat{D}_1 = 4x$ [ext \angle of Δ/<i>buite</i> \angle v Δ]</p> <p>$2x + 3x + 4x = 180^\circ$ [sum of \angles in Δ/\anglee v Δ]</p> <p>$9x = 180^\circ$</p> <p>$x = 20^\circ$</p> | <p>✓ S ✓R</p> <p>✓ S</p> <p>✓ S ✓R</p> <p>✓ answer</p> <p>(6)</p> |
| [16] | | |




QUESTION/VRAAG 9

9.1



| | | |
|------------|---|---|
| <p>9.1</p> | <p>Constr: Join BE and CD and draw h_1 from $E \perp AD$ and h_2 from $D \perp AE$</p> <p><i>Konstr: Verbind BE en CD en trek h_1 vanaf $E \perp AD$ en h_2 vanaf $D \perp AE$</i></p> <p>Proof/Bewys:</p> $\frac{\text{area } \triangle ADE}{\text{area } \triangle BDE} = \frac{\frac{1}{2} AD \times h_1}{\frac{1}{2} BD \times h_1} = \frac{AD}{BD}$ $\frac{\text{area } \triangle ADE}{\text{area } \triangle DEC} = \frac{\frac{1}{2} AE \times h_2}{\frac{1}{2} EC \times h_2} = \frac{AE}{EC}$ <p>area $\triangle ADE$ = area $\triangle ADE$ [common/gemeenskaplik]</p> <p>But area $\triangle BDE$ = area $\triangle DEC$ [same base & height ; $DE \parallel BC$ / <i>dies basis & hoogte ; $DE \parallel BC$</i>]</p> $\therefore \frac{\text{area } \triangle ADE}{\text{area } \triangle BDE} = \frac{\text{area } \triangle ADE}{\text{area } \triangle DEC}$ $\therefore \frac{AD}{BD} = \frac{AE}{EC}$ | <p>✓ constr/konstr</p> $\checkmark \frac{\text{area } \triangle ADE}{\text{area } \triangle BDE} = \frac{\frac{1}{2} AD \times h_1}{\frac{1}{2} BD \times h_1} \text{ or } \mathbf{R}$ $\checkmark \frac{\text{area } \triangle ADE}{\text{area } \triangle DEC} = \frac{AE}{EC}$ <p>✓ S ✓ R</p> <p>(6)</p> |
|------------|---|---|

9.2



| | | |
|-------|--|--|
| 9.2.1 | $\hat{A}_1 = x$ [corresp \angle s; $PQ \parallel CA$ /ooreenkomstige \angle e, $PQ \parallel CA$] $\hat{B} = x$ [\angle s opp equal sides/ \angle e teenoor gelyke sye] $\hat{A}_2 = x$ [tan-chord theorem/ \angle tussen raaklyn en koord] $\hat{P} = x$ [alt \angle s; $PQ \parallel CA$ /verw. \angle e, $PQ \parallel CA$] | \checkmark S \checkmark R \checkmark S/R \checkmark S \checkmark R \checkmark S/R |
| 9.2.2 | $\hat{B} = \hat{P}$ [proved in 9.2.1/bewys in 9.2.1] \therefore A, B, P and R are concyclic \therefore ABPR is a cyclic quadrilateral [conv \angle s in the same segment/ koord onderspan gelyke omtreks \angle e] | \checkmark S \checkmark R |
| 9.2.3 | $\frac{BA}{BQ} = \frac{BC}{BR}$ [prop th; $AC \parallel QP$] OR [line \parallel one side Δ /lyn \parallel een syn v Δ] But $QR = BR$ [sides opp = \angle s/sye teenoor = \angle e] $\therefore \frac{BA}{BQ} = \frac{BC}{QR}$ | \checkmark S \checkmark R \checkmark S |

| | | |
|-------------|---|---|
| | <p>OR</p> <p>In $\triangle ABC$ and $\triangle BQR$:</p> <p>$\hat{A}_1 = \hat{B} = x$ [proved in 9.2.1]</p> <p>$\hat{B} = \hat{Q} = x$ [proved in 9.2.1]</p> <p>$\hat{C}_1 = \hat{B}RQ = 180^\circ - 2x$ [sum of \angles of \triangle]</p> <p>$\therefore \triangle ABC \parallel \triangle BQR$</p> <p>$\therefore \frac{BA}{BQ} = \frac{BC}{QR}$</p> <p>OR</p> <p>In $\triangle ABC$ and $\triangle BQR$:</p> <p>$\hat{A}_1 = \hat{B} = x$ [proved in 9.2.1]</p> <p>$\hat{B} = \hat{Q} = x$ [proved in 9.2.1]</p> <p>$\hat{C}_1 = \hat{B}RQ = 180^\circ - 2x$ [sum of \angles of \triangle]</p> <p>$\therefore \triangle ABC \parallel \triangle BQR$ [$\angle\angle\angle$]</p> <p>$\therefore \frac{BA}{BQ} = \frac{BC}{QR}$</p> <p>OR</p> <p>In $\triangle ABC$ and $\triangle QBR$:</p> <p>\hat{B} is common</p> <p>$\hat{A}_1 = \hat{Q} = x$ [corres \angles; $PQ \parallel CA$]</p> <p>$\hat{C}_1 = \hat{B}RQ = 180^\circ - 2x$ [sum of \angles of \triangle]</p> <p>$\therefore \triangle ABC \parallel \triangle QBR$ [$\angle\angle\angle$]</p> <p>But $QR = BR$ [sides opp = \angles/sye teenoor = $\angle e$]</p> <p>$\therefore \frac{BA}{BQ} = \frac{BC}{QR}$</p> | <p>✓ S</p> <p>✓ S</p> <p>✓ S</p> <p>(3)</p> <p>✓ S</p> <p>✓ S</p> <p>✓ R</p> <p>(3)</p> <p>✓ S</p> <p>✓ S</p> <p>✓ S</p> <p>(3)</p> |
| [17] | | |

| | | |
|--------|--|---|
| 10.1.2 | <p>In $\triangle RTQ$ and $\triangle RQP$</p> <p>$\hat{T} = \hat{Q}_3$ [tan-chord theorem/<i>∠ tussen raaklyn en koord</i>]</p> <p>$\hat{Q}_1 + \hat{Q}_2 = 90^\circ$ [co-interior \angles, $MS \parallel QR$/<i>ko-binne \anglee, $MS \parallel QR$</i>] or [\angle in semi circle/<i>∠ in halwe sirkel</i>]</p> <p>$\therefore \hat{Q}_1 + \hat{Q}_2 = \hat{P} = 90^\circ$</p> <p>$\hat{R}_1 = \hat{R}_2$ [\angles of Δ/<i>∠e van Δ</i>]</p> <p>$\triangle RTQ \parallel \triangle RQP$</p> <p>$\frac{RT}{RQ} = \frac{RQ}{RP}$</p> <p>$RT = \frac{RQ^2}{RP}$</p> <p>OR</p> <p>In $\triangle RTQ$ and $\triangle RQP$</p> <p>$\hat{T} = \hat{Q}_3$ [tan-chord theorem <i>∠ tussen raaklyn en koord</i>]</p> <p>$\hat{Q}_1 + \hat{Q}_2 = 90^\circ$ [co-interior \angles, $MS \parallel QR$/<i>ko-binne \anglee, $MS \parallel QR$</i>] or [\angle in semi circle/<i>∠ in halwe sirkel</i>]</p> <p>$\therefore \hat{Q}_1 + \hat{Q}_2 = \hat{P} = 90^\circ$</p> <p>$\triangle RTQ \parallel \triangle RQP$ [\angle, \angle, \angle]</p> <p>$\frac{RT}{RQ} = \frac{RQ}{RP}$</p> <p>$RT = \frac{RQ^2}{RP}$</p> | <p>✓ S ✓ R</p> <p>✓ S</p> <p>✓ S</p> <p>✓ S</p> <p>✓ ratio</p> <p>(6)</p> <p>✓ S ✓ R</p> <p>✓ S</p> <p>✓ S</p> <p>✓ R</p> <p>✓ ratio</p> <p>(6)</p> |
| 10.2 | <p>$QR = 28$ units [midpoint theorem/<i>midpt. stelling</i>]</p> <p>$RP^2 = 28^2 - (\sqrt{640})^2$ [Pythagoras/<i>Pythagoras</i>]</p> <p>$RP = 12$ units</p> <p>$RT = \frac{RQ^2}{RP}$</p> <p>$RT = \frac{28^2}{12}$</p> <p>$RT = \frac{196}{3}$</p> <p>Radius = $\frac{98}{3}$ units</p> | <p>✓ S ✓ R</p> <p>✓ S</p> <p>✓ $RP = 12$</p> <p>✓ RT</p> <p>✓ answer</p> <p>(6)</p> |
| | | [15] |

TOTAL/TOTAAL: 150